

ORAL-PRO™

Sodium Salicylate Concentrate 48.6% w/v



FEATURES AND BENEFITS

- Aids in reducing fever caused by viral and bacterial infections.^{1 2}
- Aids in reducing fever caused by heat stress from transporting.
- Offers water solubility in hard water, and is also stable in milk.³
- Offers stress relief from vaccination and processing.
- Manufactured under cGMP standards, guaranteeing consistent quality.
- Calves: A supportive treatment of pyrexia in acute respiratory disease and in combination with appropriate (anti-infective) therapy, if necessary.

¹ Lees, P. and S.A. May, 1992. Inflammation and anti-inflammatory drugs. Pages 843–863 in: Bovine Medicine. A.H. Andrews, R.W. Blowey, H. Boyd, and R.G. Eddy, ed. Oxford Blackwell Scientific Publications, Cambridge, MA.

² Osol, A., R. Pratt, and A.R. Gennaro, 1973. The United States Dispensatory, 27th ed. J.B. Lippincott Co., Philadelphia, PA.

³ Budavari, S., M.J. O'Neil, A. Smith, and P.E. Heckelman, 1989. The Merck Index, 11th ed. Merck & Co., Rahway, NJ.

INDICATIONS

For use in the drinking water of calves, poultry and swine as an aid in reducing pain, fever and inflammation.

FOR USE IN LIVESTOCK ONLY

REORDER NO: 21004

NDC 51072-038-01

Aurora Pharmaceutical, LLC
NORTHFIELD, MINNESOTA 55057
888-215-1256 • www.aurorapharmaceutical.com

12/2011 IN 50-1108 REV 01
ss000003



MANUFACTURED IN THE USA



ORAL-PRO™

Sodium Salicylate Concentrate 48.6% w/v Solution

APPLICATION

Administer 4 ml per 100 lbs. body weight (40 mg/kg body weight Sodium Salicylate) to calves once daily for 1–3 days. Administer orally in drinking water or milk (replacer). Zero day withdrawal period for meat and offal from calves. Do not use in cows producing milk for human consumption.

INDICATIONS: For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

CALF LABEL CLAIM

Indications For Use, Calves: Supportive treatment of pyrexia in acute respiratory disease, in combination with appropriate (e.g., anti-infective) therapy if necessary.

Contraindications, Calves: Do not use Sodium Salicylate in neonates or calves less than 2 weeks of age.

Dosage, Calves: 40 mg Sodium Salicylate per kg body weight once daily, for 1–3 days. Administer orally in drinking water or milk (replacer).
Recommended withdrawal period for all food animals following use at "usual dosages": meat 24 hours.

ACTIVE INGREDIENT:

Sodium Salicylate 48.6% w/v

DIRECTIONS FOR USE:

For Analgesic and Antipyretic Use

Water Proportioner Use: Add 8 ounces (236 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

Livestock Tank Use: Add 8 ounces (236 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

For Anti-Inflammatory/Anti-Prostaglandin Use

Water Proportioner Use: Add 16 ounces (473 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Livestock Tank Use: Add 16 ounces (473 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Following Days

Water Proportioner Use: Add 10 ounces (296 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Livestock Tank Use: Add 10 ounces (296 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.
Prepare fresh solutions daily. Repeat as necessary.

WARNING: Do not administer concentrated solution by direct oral administration — gastrointestinal irritation or overdose may occur. Do not use in piglets less than 4 weeks of age. Store at controlled room temperature between 15°–30° C (59°–86° F).

CAUTION: Keep container closed when not in use. Product may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution, or place in warm water. Gently invert container to ensure uniformity of product.
Gradual darkening will not affect product stability.

REORDER NO: 21004



MANUFACTURED
IN THE USA

MANUFACTURED BY:
Aurora Pharmaceutical, LLC
NORTHFIELD, MINNESOTA 55057
888-215-1256
www.aurorapharmaceutical.com
IN 50-1108 REV 01

NDC 51072-038-01

ORAL-PRO™

Sodium Salicylate Concentrate 48.6% w/v

For Use in Livestock Only

Keep Out of Reach of Children

Net Contents:
1 Gallon (3.79 Liters)



INDICATIONS: Para uso en el agua potable de las aves de corral y cerdos como una ayuda para reducir el dolor, fiebre e inflamación.

TERNEROS DE LA ETIQUETA

Indicaciones de Uso, Terneros: Tratamiento de la fiebre de enfermedad respiratoria aguda, en combinación con la adecuada (por ejemplo, anti-infecciosa), la terapia, si es necesario.

Contraindicaciones, Terneros: No utilice salicilato de sodio en los recién nacidos o terneros de menos de 2 semanas de edad.

Dosis, Terneros: 40 mg de salicilato sódico por kg de peso corporal una vez al día, durante 1–3 días. Administrar por vía oral en el agua potable o la leche (sustituto). De suspensión recomendados período para todos los animales de alimentos después de su uso en "dosis habituales": la carne de 24 horas.

INGREDIENTES ACTIVOS

Salicilato de Sodio 48.6% p/b

MODO DE EMPLEO:

Para el Uso de Analgésicos y Antipiréticos

Uso del Dosificador de Agua: Agregue 8 onzas (236 ml) de concentrado de salicilato de sodio 48.6% para hacer un galón de solución madre y administre a través de un contenedor medido a 1:128 (1 onza por galón). Esto permitirá alcanzar una dosis objetivo de 11.3 mg/lb (25 mg/kg) de peso corporal al día.

Uso del Tanque Ganadería: Agregue 8 onzas (236 ml) de concentrado de salicilato de sodio 48.6% a 128 galones de agua potable. Esto permitirá alcanzar una dosis objetivo de 11.3 mg/lb (25 mg/kg) de peso corporal al día.

Para el Uso Anti-Inflamatorio/Anti-Prostaglandina
Uso del Dosificador de Agua: Agregue 16 onzas (473 ml) de concentrado de salicilato de sodio 48.6% para hacer un galón de solución madre y administre a través de un contenedor medido a 1:128 (1 onza por galón). Esto permitirá alcanzar una dosis objetivo de 22.7 mg/lb (50 mg/kg) de peso corporal al día.

Uso del Tanque Ganadería: Agregue 16 onzas (473 ml) de concentrado de salicilato de sodio 48.6% a 128 galones de agua potable. Esto permitirá alcanzar una dosis objetivo de 22.7 mg/lb (50 mg/kg) de peso corporal al día.

Los Días Siguyentes

Uso del Dosificador de Agua: Agregue 10 onzas (296 ml) de concentrado de salicilato de sodio 48.6% para hacer un galón de solución madre y administre a través de un contenedor medido a 1:128 (1 onza por galón). Esto permitirá alcanzar una dosis objetivo de 13.6 mg/lb (30 mg/kg) de peso corporal al día.

Uso del Tanque Ganadería: Agregue 10 onzas (296 ml) de concentrado de salicilato de sodio 48.6% a 128 galones de agua potable. Esto permitirá alcanzar una dosis objetivo de 13.6 mg/lb (30 mg/kg) de peso corporal al día.
Prepare soluciones frescas al día. Repita según sea necesario.

ADVERTENCIA: No administrar la solución concentrada por vía oral en directo — irritación gastrointestinal o una sobredosis puedan ocurrir. No utilizar en lechones de menos de 4 semanas de edad.
Conservar a temperatura ambiente controlada entre 15°–30° C (59°–86° F).

PRECAUCIÓN: Mantenga el recipiente cerrado cuando no este en uso. El producto puede solidificarse a temperaturas de almacenamiento en frío. Coloque el recipiente en el almacenamiento a temperatura ambiente, el cual se derretirá en la solución, o en lugar con agua tibia. Invierta suavemente el recipiente para asegurar la uniformidad del producto.
Oscurecimiento gradual no afectará a la estabilidad del producto.



ABOUT AURORA PHARMACEUTICAL

Aurora is devoted to improved formulations and delivery systems that are consumer friendly and which utilize state-of-the-art ingredient "green" technologies. We are committed to manufacturing products with top-quality ingredients that increase the stability, shelf life and efficacy of our products while minimizing harmful ingredients and environmental contamination.

Aurora Pharmaceutical, LLC
NORTHFIELD, MINNESOTA 55057
888-215-1256 • www.aurorapharmaceutical.com



MANUFACTURED IN THE USA