

# ORAL-PRO™

## Sodium Salicylate Concentrate 60% w/v with Caffeine 5.7% w/v



### FEATURES AND BENEFITS

- 60% Sodium Salicylate Concentrate with Caffeine for use in the drinking water of swine and poultry. Caffeine added at 9.6% of sodium salicylate content.
- Aids in reducing fever caused by viral and bacterial infections.<sup>1 2</sup>
- Aids in reducing fever caused by heat stress from transporting.
- Offers water solubility in hard water, and is also stable in milk.<sup>3</sup>
- Offers stress relief from vaccination and processing.
- Manufactured under cGMP standards, guaranteeing consistent quality.

<sup>1</sup> Lees, P. and S.A. May, 1992. Inflammation and anti-inflammatory drugs. Pages 843–863 in: Bovine Medicine. A.H. Andrews, R.W. Blowey, H. Boyd, and R.G. Eddy, ed. Oxford Blackwell Scientific Publications, Cambridge, MA.

<sup>2</sup> Osol, A., R. Pratt, and A.R. Gennaro, 1973. The United States Dispensatory, 27th ed. J.B. Lippincott Co., Philadelphia, PA.

<sup>3</sup> Budavari, S., M.J. O'Neil, A. Smith, and P.E. Heckelman, 1989. The Merck Index, 11th ed. Merck & Co., Rahway, NJ.

### INDICATIONS

For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

### FOR USE IN LIVESTOCK ONLY

REORDER NO: 21003

NDC 51072-039-01

**Aurora Pharmaceutical, LLC**  
NORTHFIELD, MINNESOTA 55057  
888-215-1256 • [www.aurorapharmaceutical.com](http://www.aurorapharmaceutical.com)

12/2011 IN 50-1108 REV 01  
ss000002



MANUFACTURED IN THE USA



# ORAL-PRO™

## Sodium Salicylate Concentrate 60% w/v Solution with Caffeine 5.7% w/v

### ABOUT AURORA PHARMACEUTICAL

Aurora is devoted to improved formulations and delivery systems that are consumer friendly and which utilize state-of-the-art ingredient “green” technologies. We are committed to manufacturing products with top-quality ingredients that increase the stability, shelf life and efficacy of our products while minimizing harmful ingredients and environmental contamination.

**INDICATIONS:** For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

**ACTIVE INGREDIENTS:**  
Sodium Salicylate . . . . . 60% w/v      Caffeine . . . . . 5.7% w/v

**DIRECTIONS FOR USE:**

**For Analgesic and Antipyretic Use**

**Water Proportioner Use:** Add 6.5 ounces (192 ml) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

**Livestock Tank Use:** Add 6.5 ounces (192 ml) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

**For Anti-inflammatory/Anti-Prostaglandin Use**

**Water Proportioner Use:** Add 14 ounces (414 ml) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

**Livestock Tank Use:** Add 14 ounces (414 ml) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

**Following Days**

**Water Proportioner Use:** Add 8 ounces (236 ml) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

**Livestock Tank Use:** Add 8 ounces (236 ml) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Prepare fresh solutions daily. Repeat as necessary.

**WARNING:** Do not administer concentrated solution by direct oral administration — gastrointestinal irritation or overdose may occur. Do not use in piglets less than 4 weeks of age.

Store at controlled room temperature between 15°–30° C (59°–86° F).

**CAUTION:** Keep container closed when not in use. Product may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution, or place in warm water. Gently invert container to ensure uniformity of product.

Gradual darkening will not affect product stability.

REORDER NO: 21003



MANUFACTURED  
IN THE USA

MANUFACTURED BY:  
**Aurora Pharmaceutical, LLC**  
NORTHFIELD, MINNESOTA 55057  
888-215-1256  
[www.aurorapharmaceutical.com](http://www.aurorapharmaceutical.com)  
IN 50-1110 REV 01

NDC 51072-039-01

# ORAL-PRO™

## Sodium Salicylate Concentrate 60% w/v with Caffeine 5.7% w/v

For Use in Livestock Only

Keep Out of Reach of Children

Net Contents:  
**1 Gallon (3.79 Liters)**



**INDICACIONES:** Para uso en el agua potable de las aves de corral y cerdos como una ayuda para reducir el dolor, fiebre e inflamación.

**INGREDIENTES ACTIVOS**  
Salicilato de Sodio . . . . . 60% p/b      Caffeína . . . . . 5.7% p/b

**MODO DE EMPLEO:**

**Para el Uso de Analgésicos y Antipiréticos**

**Uso del Dosificador de Agua:** Agregue 6.5 onzas (192 ml) de concentrado de salicilato de sodio 60% con cafeína 5.7% para hacer un galón de solución madre y administre a través de un contenedor medido a 1:128 (1 onza por galón). Esto permitirá alcanzar una dosis objetivo de 11.3 mg/lb (25 mg/kg) de peso corporal al día.

**Uso del Tanque Ganadería:** Agregue 6.5 onzas (192 ml) de concentrado de salicilato de sodio concentrado 60% con cafeína 5.7% a 128 galones de agua potable. Esto permitirá alcanzar una dosis objetivo de 11.3 mg/lb (25 mg/kg) de peso corporal al día.

**Para el Uso Anti-inflamatorio/Anti-Prostaglandina**

**Uso del Dosificador de Agua:** Agregue 14 oz (414 ml) de concentrado de salicilato de sodio 60% con cafeína 5.7% para hacer un galón de solución madre y administre a través de un contenedor medido a 1:128 (1 onza por galón). Esto permitirá alcanzar una dosis objetivo de 22.7 mg/lb (50 mg/kg) de peso corporal al día.

**Uso del Tanque Ganadería:** Agregue 14 oz (414 ml) de concentrado de salicilato de sodio 60% con cafeína 5.7% a 128 galones de agua potable. Esto permitirá alcanzar una dosis objetivo de 22.7 mg/lb (50 mg/kg) de peso corporal al día.

**Los Días Siguyentes**

**Uso del Dosificador de Agua:** Agregue 8 onzas (236 ml) de concentrado de salicilato de sodio 60% con cafeína 5.7% para hacer un galón de solución madre y administre a través de un contenedor medido a 1:128 (1 onza por galón). Esto permitirá alcanzar una dosis objetivo de 13.6 mg/lb (30 mg/kg) de peso corporal al día.

**Uso del Tanque Ganadería:** Agregue 8 onzas (236 ml) de concentrado de salicilato de sodio 60% con cafeína 5.7% a 128 galones de agua potable. Esto permitirá alcanzar una dosis objetivo de 13.6 mg/lb (30 mg/kg) de peso corporal al día.

Prepare soluciones frescas al día. Repita según sea necesario.

**ADVERTENCIA:** No administrar la solución concentrada por vía oral en directo — irritación gastrointestinal o una sobredosis pueden ocurrir. No utilizar en lechones de menos de 4 semanas de edad.

Conserve a temperatura ambiente controlada entre 15°–30° C (59°–86° F).

**PRECAUCIÓN:** Mantenga el recipiente cerrado cuando no este en uso. El producto puede solidificarse a temperaturas de almacenamiento en frío. Coloque el recipiente en el almacenamiento a temperatura ambiente, el cual se derretirá en la solución, o en lugar con agua tibia. Invierta suavemente el recipiente para asegurar la uniformidad del producto. Oscurecimiento gradual no afectará a la estabilidad del producto.

