

ORAL-PRO™

Sodium Salicylate Concentrate 48.6% w/v



FEATURES AND BENEFITS

- Aids in reducing fever caused by viral and bacterial infections.^{1 2}
- Aids in reducing fever caused by heat stress from transporting.
- Offers water solubility in hard water, and is also stable in milk.³
- Offers stress relief from vaccination and processing.
- Manufactured under cGMP standards, guaranteeing consistent quality.
- Calves: A supportive treatment of pyrexia in acute respiratory disease and in combination with appropriate (anti-infective) therapy, if necessary.

¹ Lees, P. and S.A. May, 1992. Inflammation and anti-inflammatory drugs. Pages 843–863 in: Bovine Medicine. A.H. Andrews, R.W. Blowey, H. Boyd, and R.G. Eddy, ed. Oxford Blackwell Scientific Publications, Cambridge, MA.

² Osol, A., R. Pratt, and A.R. Gennaro, 1973. The United States Dispensatory, 27th ed. J.B. Lippincott Co., Philadelphia, PA.

³ Budavari, S., M.J. O'Neil, A. Smith, and P.E. Heckelman, 1989. The Merck Index, 11th ed. Merck & Co., Rahway, NJ.

INDICATIONS

For use in the drinking water of calves, poultry and swine as an aid in reducing pain, fever and inflammation.

FOR USE IN LIVESTOCK ONLY

REORDER NO: 21004

NDC 51072-038-01

Aurora Pharmaceutical, LLC
NORTHFIELD, MINNESOTA 55057
888-215-1256 • www.aurorapharmaceutical.com

06/2016 / IN 50-1108 06/2016
ss000003



MANUFACTURED IN THE USA



ORAL-PRO™

Sodium Salicylate Concentrate 48.6% w/v Solution

APPLICATION

Administer 4 mL per 100 lbs. body weight (40 mg/kg body weight Sodium Salicylate) to calves once daily for 1–3 days. Administer orally in drinking water or milk (replacer). Zero day withdrawal period for meat and offal from calves. Do not use in cows producing milk for human consumption.

ACTIVE INGREDIENT
Sodium Salicylate 48.6% w/v

CALF LABEL CLAIM
Indications For Use, Calves: Supportive treatment of pyrexia in acute respiratory disease, in combination with appropriate (e.g., anti-infective) therapy if necessary.
Contraindications, Calves: Do not use Sodium Salicylate in neonates or calves less than 2 weeks of age.
Dosage, Calves: 40 mg Sodium Salicylate per kg body weight (4 mL per 100 lbs.) once daily, for 1–3 days. Administer orally in drinking water or milk (replacer).

SWINE AND POULTRY LABEL CLAIM
Indications: For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

DIRECTIONS FOR USE
For Analgesic and Antipyretic Use
Water Proportioner Use: Add 8 ounces (236 mL) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.
Livestock Tank Use: Add 8 ounces (236 mL) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

For Anti-inflammatory/Anti-Prostaglandin Use — Day 1
Water Proportioner Use: Add 16 ounces (473 mL) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

MANUFACTURED IN THE USA

REORDER NO: 21004
MANUFACTURED BY:
Aurora Pharmaceutical, LLC
NORTHFIELD, MINNESOTA 55057
888-215-1256
www.aurorapharmaceutical.com
IN 50-1108 06/2016

NDC 51072-038-01



Sodium Salicylate Concentrate 48.6% w/v

For Use in Livestock Only

Keep Out of Reach of Children

Net Contents:
1 Gallon (3.79 Liters)



DIRECTIONS FOR USE (continued)

Livestock Tank Use: Add 16 ounces (473 mL) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Day 2 through 7
Water Proportioner Use: Add 10 ounces (296 mL) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.
Livestock Tank Use: Add 10 ounces (296 mL) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily. Prepare fresh solutions daily. Repeat as necessary.

WARNING
Do not administer concentrated solution by direct oral administration — gastro-intestinal irritation or overdose may occur. Do not use in piglets less than 3 weeks of age.
Store at 20°–25° C (68°–77° F). Excursions permitted between 15°–30° C (59°–86° F).
Do not use if allergic or sensitive to the active ingredients.
In California, Restricted Drug — Use Only as Directed.

CAUTION
Keep container closed when not in use. Product may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution, or place in warm water. Gently invert container to ensure uniformity of product. Gradual darkening will not affect product stability.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away.

TAKE TIME  OBSERVE LABEL DIRECTIONS



ABOUT AURORA PHARMACEUTICAL

Aurora is devoted to improved formulations and delivery systems that are consumer friendly and which utilize state-of-the-art ingredient “green” technologies. We are committed to manufacturing products with top-quality ingredients that increase the stability, shelf life and efficacy of our products while minimizing harmful ingredients and environmental contamination.