

ORAL-PRO™

Sodium Salicylate Concentrate 60% w/v with Caffeine 5.7% w/v



FEATURES AND BENEFITS

- For use in the drinking water of swine and poultry. Caffeine added at 9.6% of sodium salicylate content.
- Aids in reducing fever caused by viral and bacterial infections.^{1 2}
- Aids in reducing fever caused by heat stress from transporting.
- Offers water solubility in hard water, and is also stable in milk.³
- Offers stress relief from vaccination and processing.
- Manufactured under cGMP standards, guaranteeing consistent quality.

¹ Lees, P. and S.A. May, 1992. Inflammation and anti-inflammatory drugs. Pages 843–863 in: Bovine Medicine. A.H. Andrews, R.W. Blowey, H. Boyd, and R.G. Eddy, ed. Oxford Blackwell Scientific Publications, Cambridge, MA.

² Osol, A., R. Pratt, and A.R. Gennaro, 1973. The United States Dispensatory, 27th ed. J.B. Lippincott Co., Philadelphia, PA.

³ Budavari, S., M.J. O'Neil, A. Smith, and P.E. Heckelman, 1989. The Merck Index, 11th ed. Merck & Co., Rahway, NJ.

INDICATIONS

For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

FOR USE IN LIVESTOCK ONLY

REORDER NO: 21003

NDC 51072-039-01

Aurora Pharmaceutical, LLC
NORTHFIELD, MINNESOTA 55057
888-215-1256 • www.aurorapharmaceutical.com

05/2016 / IN 50-1110 05/2016
ss000002



MANUFACTURED IN THE USA



ORAL-PRO™

Sodium Salicylate Concentrate 60% w/v Solution with Caffeine 5.7% w/v

ABOUT AURORA PHARMACEUTICAL

Aurora is devoted to improved formulations and delivery systems that are consumer friendly and which utilize state-of-the-art ingredient “green” technologies. We are committed to manufacturing products with top-quality ingredients that increase the stability, shelf life and efficacy of our products while minimizing harmful ingredients and environmental contamination.

ACTIVE INGREDIENTS

Sodium Salicylate60% w/v
Caffeine5.7% w/v

NDC 51072-039-01

INDICATIONS: For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

DIRECTIONS FOR USE:

For Analgesic and Antipyretic Use

Water Proportioner Use: Add 6.5 ounces (192 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

Livestock Tank Use: Add 6.5 ounces (192 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

For Anti-Inflammatory/Anti-Prostaglandin Use — Day 1

Water Proportioner Use: Add 14 ounces (414 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Livestock Tank Use: Add 14 ounces (414 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

MANUFACTURED IN THE USA



REORDER NO: 21003

MANUFACTURED BY:
Aurora Pharmaceutical, LLC
NORTHFIELD, MINNESOTA 55057
888-215-1256
www.aurorapharmaceutical.com
IN 50-1110 05/2016

ORAL-PRO™

Sodium Salicylate Concentrate 60% w/v with Caffeine 5.7% w/v

For Use in Livestock Only

Keep Out of Reach of Children

Net Contents:
1 Gallon (3.79 Liters)



DIRECTIONS FOR USE (continued)

Day 2 through 7

Water Proportioner Use: Add 8 ounces (236 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Livestock Tank Use: Add 8 ounces (236 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.
Prepare fresh solutions daily. Repeat as necessary.

WARNING

Do not administer concentrated solution by direct oral administration — gastro-intestinal irritation or overdose may occur. Do not use in piglets less than 3 weeks of age.

Store at 20°–25° C (68°–77° F). Excursions permitted between 15°–30° C (59°–86° F).

Do not use if allergic or sensitive to the active ingredients.

In California, Restricted Drug — Use Only as Directed.



CAUTION

Keep container closed when not in use. Product may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution, or place in warm water. Gently invert container to ensure uniformity of product.
Gradual darkening will not affect product stability.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

