



Wound Care Spray With Pain Relief 2% Available Iodine With Lidocaine

INDICATIONS

For topical use in animals as an aid in reducing pain, licking and chewing on wounds and surgical incisions. Kills up to 99% of surface germs in 15 seconds or less that can potentially cause an infection.

- A 2% titratable iodine that kills up to 99% of surface germs that may cause an infection
- Contains lidocaine to alleviate pain in wounds and surgical incisions
- Dries in approximately 60 seconds and forms a highly visible and durable coating
- Has a natural anti-biting/chewing agent to help prevent self-mutilation and/or cohort licking, biting and chewing



AVAILABILITY

16 oz. Bottle - Reorder No: 19001 1 Gallon - Reorder No: 19002



The 16 oz. bottle comes with a trigger sprayer. For refills, pour the 1 gallon into the spray bottle dialed to a spray pattern. Hold sprayer about 4—6 inches from area to be treated. Spray 1 or more times - allow to dry. Repeat as needed. Forms a highly visible coating that is durable.



WARNINGS

Do not use in the eyes. Do not use if you are allergic to povidone-iodine. Prolonged exposure may cause irritation. If swallowed, get medical help or contact a Poison Control Center right away.

Store at 20°-25° C (68°-77° F). Excursions permitted between 15°-30° C (59°-86° F). Avoid inhalation.

CAUTION

Flammable - use with caution. For external use only. Keep out of reach of children. Wet hair or fur is flammable.

FEATURE	BARRIER® WOUND CARE SPRAY	TRIODINE-7®	IODINE 1% SOLUTION	STOP BITE™
Labeled for Animals				
nstant Kill < 10 sec.				
Prolonged Kill > 10 sec.				
Pain Relief				
Anti-Chew/Anti-Bite				
Non-Staining				
Scented				
Fast Drying				
Washes Off Hands				
Visible				
Residual Activity		THE RESERVE OF THE PARTY OF THE		

