# ORAL-PRO



Sodium Salicylate Concentrate 60% w/v with Caffeine 5.7% w/v

#### **INDICATIONS**

For use in the drinking water of poultry as an aid in reducing pain, fever and inflammation.

- Caffeine added at 9.6% of sodium salicylate content
- Aids in reducing fever caused by viral and bacterial infections<sup>1,2</sup>
- Aids in reducing fever caused by heat stress from transporting
- Offers water solubility in hard water
- Offers stress relief from vaccination and processing
- Manufactured under cGMP standards, guaranteeing consistent quality



#### **AVAILABILITY**

1 Gallon - Reorder No: 21003

1 Lees, P. and S.A. May, 1992. Inflammation and anti-inflammatory drugs. Pages 843–863 in: Bovine Medicine. A.H. Andrews, R.W. Blowey, H. Boyd, and R.G. Eddy, ed. Oxford Blackwell Scientific Publications, Cambridge, MA. 2 Osol, A., R. Pratt, and A.R. Gennaro, 1973. The United States Dispensatory, 27th ed. J.B. Lippincott Co., Philadelphia, PA.



NDC 51072-039-01

### ORAL-PRO®

Sodium Salicylate Concentrate 60% w/v

with Caffeine 5.7% w/v

For Use in Livestock Only

#### **Keep Out of Reach of Children**

1 Gallon (3.79 L)

#### **ACTIVE INGREDIENTS**

Sodium Salicylate.....60% w/v Caffeine .....5.7% w/v

**INDICATIONS:** For use in the drinking water of poultry as an aid in reducing pain, fever and inflammation.

#### **DIRECTIONS FOR USE:**

For Analgesic and Antipyretic Use

Water Proportioner Use: Add 6.5 ounces (192 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

Livestock Tank Use: Add 6.5 ounces (192 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

## For Anti-Inflammatory/Anti Prostaglandin Use Day 1

Water Proportioner Use: Add 14 ounces (414 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Livestock Tank Use: Add 14 ounces (414 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

#### Day 2 Through 7

Water Proportioner Use: Add 8 ounces (236 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Livestock Tank Use: Add 8 ounces (236 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily. Prepare fresh solutions daily. Repeat as necessary.

#### WARNING

Do not administer concentrated solution by direct oral administration - gastrointestinal irritation or overdose may occur. Do not use in piglets less than 3 weeks of age.

Store at  $20^{\circ}$ – $25^{\circ}$  C ( $68^{\circ}$ – $77^{\circ}$  f). Excursions permitted between  $15^{\circ}$ – $30^{\circ}$  C ( $59^{\circ}$ – $86^{\circ}$  f).

Do not use if allergic or sensitive to the active ingredients.

#### **CAUTION**

Keep container closed when not in use. Product may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution, or place in warm water. Gently invert container to ensure uniformity of product. Gradual darkening will not affect product stability.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.



